

Research

Outcomes of planned home birth with registered midwife versus planned hospital birth with midwife or physician

Patricia A. Janssen, PhD, Lee Saxell, MA, Lesley A. Page, PhD, Michael C. Klein, MD, Robert M. Liston, MD and Shoo K. Lee, MBBS PhD

From the School of Population and Public Health (Janssen), the Departments of Family Practice (Klein) and Obstetrics and Gynecology (Janssen, Liston) and the Division of Midwifery (Saxell), Faculty of Medicine, University of British Columbia, Vancouver, BC; the Child and Family Research Institute (Janssen, Klein, Liston), Vancouver, BC; the Nightingale School of Nursing and Midwifery (Page), King's College, London, UK; the Department of Pediatrics (Lee); and the Integrated Centre for Care Advancement Through Research (Lee), University of Alberta, Edmonton, Alta.

Correspondence to: Dr. Patricia A. Janssen, School of Population and Public Health, University of British Columbia, 5804 Fairview Cres., Vancouver BC V6T 1Z3; fax 604 806-8006; pjanssen@interchange.ubc.ca

Background: Studies of planned home births attended by registered midwives have been limited by incomplete data, nonrepresentative sampling, inadequate statistical power and the inability to exclude unplanned home births. We compared the outcomes of planned home births attended by midwives with those of planned hospital births attended by midwives or physicians.

This Article

- ▶ [Figures Only](#)
- ▶ [Full Text](#)
- ▶ [Full Text \(PDF\)](#)
- ▶ [Online Appendix](#)
- ▶ **All Versions of this Article:**
[cmaj.081869v1](#)
181/6-7/377 *most recent*
- ▶ [Submit a response](#)
- ▶ [View responses](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me when eLetters are posted](#)
- ▶ [Alert me if a correction is posted](#)

Services

- ▶ [Email this article to a friend](#)
- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)

Google Scholar

- ▶ [Articles by Janssen, P. A.](#)
- ▶ [Articles by Lee, S. K.](#)

PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Janssen, P. A.](#)
- ▶ [Articles by Lee, S. K.](#)

Related Collections

- ▶ [Other obstetrics & gynecology](#)
- ▶ [Related Articles](#)

of planned home births attended by midwives with those of planned hospital births attended by midwives or physicians.

Methods: We included all planned home births attended by registered midwives from Jan. 1, 2000, to Dec. 31, 2004, in British Columbia, Canada ($n = 2889$), and all planned hospital births meeting the eligibility requirements for home birth that were attended by the same cohort of midwives ($n = 4752$). We also included a matched sample of physician-attended planned hospital births ($n = 5331$). The primary outcome measure was perinatal mortality; secondary outcomes were obstetric interventions and adverse maternal and neonatal outcomes.

Results: The rate of perinatal death per 1000 births was 0.35 (95% confidence interval [CI] 0.00–1.03) in the group of planned home births; the rate in the group of planned hospital births was 0.57 (95% CI 0.00–1.43) among women attended by a midwife and 0.64 (95% CI 0.00–1.56) among those attended by a physician. Women in the planned home-birth group were significantly less likely than those who planned a midwife-attended hospital birth to have obstetric interventions (e.g., electronic fetal monitoring, relative risk [RR] 0.32, 95% CI 0.29–0.36; assisted vaginal delivery, RR 0.41, 95% CI 0.33–0.52) or adverse maternal outcomes (e.g., third- or fourth-degree perineal tear, RR 0.41, 95% CI 0.28–0.59; postpartum hemorrhage, RR 0.62, 95% CI 0.49–0.77). The findings were similar in the comparison with physician-assisted hospital births. Newborns in the home-birth group were less likely than those in the midwife-attended hospital-birth group to require resuscitation at birth (RR 0.23, 95% CI 0.14–0.37) or oxygen therapy beyond 24 hours (RR 0.37, 95% CI 0.24–0.59). The findings were similar in the comparison with newborns in the physician-assisted hospital births; in addition, newborns in the home-birth group were less likely to have meconium aspiration (RR 0.45, 95% CI 0.21–0.93) and more likely to be admitted to hospital or readmitted if born in hospital (RR 1.39, 95% CI 1.09–1.85).

Interpretation: Planned home birth attended by a registered midwife was associated with very low and comparable rates of perinatal death and reduced rates of obstetric interventions and other adverse perinatal outcomes compared with planned hospital birth attended by a midwife or physician.

Related Articles

Highlights

Can. Med. Assoc. J. 2009 181: 353. [\[Full Text\]](#) [\[PDF\]](#)

Dans ce numéro

Can. Med. Assoc. J. 2009 181: E101. [\[Full Text\]](#) [\[PDF\]](#)

The safety of home birth: Is the evidence good enough?

Helen McLachlan and Della Forster

Can. Med. Assoc. J. 2009 181: 359-360. [\[Full Text\]](#) [\[PDF\]](#)

eLetters:

Read all [eLetters](#)

Why include stillbirths before 28 weeks?

Amy B. Tuteur, MD

cmaj.ca, 2 Sep 2009 [\[Full text\]](#)

Re: Why include stillbirths before 28 weeks?

Lee Saxell

cmaj.ca, 8 Sep 2009 [\[Full text\]](#)

[HOME](#) [CURRENT ISSUE](#) [PAST ISSUES](#) [COLLECTIONS](#) [HELP](#) [SEARCH](#)

Copyright 1995-2009, Canadian Medical Association. All rights reserved. ISSN 1488-2329 (e) 0820-3946
(p)

All editorial matter in CMAJ represents the opinions of the authors and not necessarily those of the Canadian Medical Association.